# STEPHAN J. SWEET, M.D., M.P.H

A Professional Corporation Sports Medicine, Arthroscopy & General Orthopedic Surgery

## POST OPERATIVE INSTRUCTION-ARTHSCOPIC SHOULDER SURGERY-ACROMIOPLASTY/ AC RESECTION

#### **WOUND CARE**

- You may remove the Operative Dressing on Post- Op Day #2.
- Apply Band-Aids to the wounds. Change the Band-Aids daily until the wounds are completely dry. Please do not use Bacitracin or other ointments under the bandage.
- Use Ice or cold therapy device 30-40 minutes 3-4 times per day until swelling is down, then as needed for pain relief.
- You may shower on Post- Op Day #2. Please cover the wound with plastic wrap and secure it to your skin
  with tape. You may remove the sling for showering, but keep the arm across the chest. Gently pat the
  area dry. <u>Do not soak</u> the shoulder in water. <u>Do not go swimming</u> in the pool or ocean until clear by Dr.
  Sweet.

### **EXERCISES**

- Wear the sling at all times except when doing your exercises. You may remove the sling for showering, but keep the arm across the chest. When your shoulder begins to feel comfortable out of the sling, you may remove it (3-7 days).
- Please perform the exercises on the attached home Exercise Program 2-3 times daily:
  - Pendulums and shoulder range of motion
  - Elbow/ Hand/ Wrist Range of Motion Exercises
  - Grip Strengthening
  - Scapular Tightening

## **POSTOP**

- You will be given three prescriptions for pain relief for you to use post-operatively:
  - NAPROSYN, an anti-inflammatory medication, is to be used twice a day with breakfast and dinner, for the first seven days after surgery
    - \*\* IF YOU HAVE HISTORY OF AN ULCER OR DIFFICULTY USING ANTI-INFLAMMATORIES OR ASPIRIN >> DO NOT USE
  - Norco <u>or Percocet</u>, a strong narcotic, to be used only on an "<u>as needed"</u> basis for pain in addition to the Naprosyn
  - Colace, a stool softener if you're experiencing constipation.
- If you have any adverse effects with the medications, please call our office.
- If you develop a Fever (101.5), Redness or Drainage from the surgical incision site, please call our office to arrange for an evaluation.

### **FOLLOW-UP**

Please call the office to schedule a follow up appointment for your wound check 10-14 days post
operatively if it has not already been scheduled. Dr. Sweet will prescribe physical therapy at this
time.

### IF YOU HAVE ANY QUESTIONS, PLEASE FEEL FREE TO CALL OUR OFFICE